

Food and Feeding Habits of Grey Mulletts (Pisces: Mugilidae) in Two Estuaries in Ghana

H. R. Dankwa^{1*}, J. Blay Jr², K. Yankson²

¹CSIR-Water Research Institute, P. O. Box 38, Achimota, Ghana

²University of Cape Coast, Cape Coast, Ghana

*Corresponding author, E-mail: hrdankwa@hotmail.com

Abstract

Food and feeding habits of grey mulletts (Mugilidae) in the River Volta and River Pra estuaries in Ghana were studied between February 1997 and July 1998 as part of efforts to encourage their culture. Stomach contents of fish samples, obtained with a cast net and a drag net, were analysed using the 'points' and frequency of occurrence method. Diatoms, detrital material and sand particles were the major items in the stomachs of all the species from the two estuaries. Their diet did not show any substantial seasonality, neither did it change with size. The various species ingested sand particles of selected range with *Liza dumerilii* ingesting the widest range in both estuaries, 41.2-1195.8 μm in the Volta estuary, and 33.0-1649.0 μm in the Pra estuary. Species that ingested the same modal size of sand particles showed preferences for different food items. The shortest mean relative gut length (gut length to body length ratio) of 1.82 and 1.72 were calculated for *L. dumerilii* in the Volta and Pra estuaries, respectively, while the longest mean relative gut length of 4.56 was calculated for *Mugil cephalus* in the Volta estuary and 4.33 for *Liza grandisquamis* in the Pra estuary. All the species showed a diurnal feeding habit, with the main feeding period occurring between 08.00 and 12.00 h. The peak feeding time, however, differed among the species.

Introduction

Grey mulletts contribute substantially to the fishery of tropical and subtropical regions of the world (Wijeyaratne & Costa, 1986; Koutrakis, Sinis & Economidis, 1994). Subsequently, they are a major source of subsistence protein requirements of the peoples of many countries (ICLARM, 1980), including Ghana. Their characteristics, including the ability to grow to a large size and ready availability, have made the group an excellent choice for culture (Bardach, Ryther & McLaren, 1972). For a successful culture operation of any species, however, a good understanding of the biology of the species is an important pre-requisite. Food and feeding habits of grey mulletts have, therefore, been a major research area (Marais, 1980; Ferrari & Chierigato, 1981; Albaret & Legendre, 1985; Blaber, 1987; King, 1988; Ikomi, 1990).

The only work done in Ghana on food and feeding habits of grey mulletts is by Blay (1995a,b) in the Elmina and Cape Coast lagoons. Thus, information on the food and feeding habits of grey mulletts in Ghana, especially from estuarine ecosystems and the offshore marine waters, is limited. This study was undertaken as part of efforts to encourage the culture of grey mulletts in Ghana to enhance local fish production.

Materials and methods

The study was conducted at the estuaries of River Pra and River Volta, both of which discharge into the Gulf of Guinea (Fig. 1). The River Volta estuary lies within the coastal savanna zone with an annual rainfall of 750–1,250 mm (Dickson & Benneh, 1977). The estuary is about 1.2 km wide at the mouth. Two dams have been built on the River Volta at Akosombo and further down at Kpong, and these, to a large extent, influence the quality and quantity of water at the estuary.




Fig. 1. Map of Ghana showing the location of the Volta and Pra estuaries

The River Pra basin lies in the moist evergreen forest zone of Ghana, with an annual rainfall of between 1,500 and 1,750 mm (Dickson & Benneh, 1977). Its banks, unlike the Volta estuary, where most of the banks are settlements, are fringed by red mangroves (*Rhizophora* spp.) up to about 10 km inland (Obodai, Yankson & Blay Jr, 1996). The estuary is approximately 100 m wide at the point of entry into the sea and the basin is subjected to impacts of mining activities upriver.

Grey mullets were sampled from the estuaries each month from February 1997 to July 1998. The fish were caught with a cast net and a drag net. Samples were also obtained from the catches of local fishermen to augment the experimental catches. Two 24-h samplings were undertaken during the periods 11–12 November 1998 and 9–10 June 1999 in the Volta estuary to determine the daily feeding periodicity. Sampling was done at four-hourly intervals with a seine net. Two hauls were made at each sampling time, both of which lasted about 80 min. Conditions in the Pra estuary precluded the use of a dragnet, which was the only means of collecting sufficient samples for studies on the daily feeding cycle.

Mullet samples were kept in 10% formalin and later sorted out into the various species using identification keys (Schneider, 1990; Lévêque, Paugy & Teugels, 1992). The total length (TL) and standard length (SL) of the fish were measured to the nearest 1.0 mm and weighed to the nearest 0.1 g. The stomach contents of each individual were also

weighed to the nearest 0.01 g and the total length of the intestine measured to the nearest 1.0 mm to calculate the relative gut length.

In each month, between five and 10 stomachs of each species were preserved in 5% formalin. Points were awarded each stomach according to its degree of fullness using an arbitrary 4-point scale (20, 15, 10 and 5 points for full, $\frac{3}{4}$, $\frac{1}{2}$ and $\frac{1}{4}$ filled stomach, respectively). The food items were identified under high power objectives of the microscope. Further analysis of the stomach contents was carried out to assess the percentage composition and frequency of occurrence of the food items (Hynes, 1950).

The stomach contents were broadly grouped into polychaetes, zooplankton, diatoms, green algae, blue-green algae, detritus and sand particles. Using a calibrated eyepiece micrometer, random measurements were made of the longest axis of sand particles from stomachs of the grey mullet species to determine particle size preference among species and between sites. The Stomach Fullness Index (FI) was used as an indicator of feeding intensity to evaluate seasonal feeding activity as well as diel feeding periodicity of the various species. It was calculated using the equation:

$$FI = (\text{Weight of stomach contents} \times 100) / \text{Fish weight (Hureau, 1966)}.$$

Samples used for the calculation of FI were those captured between 8.00 h and 11.00 h since that was the most active period for feeding. A cluster analysis was performed to illustrate similarities among the species based on the frequency of stomach contents.

Results

Size range

The size range of the different species of grey mullets in the two estuaries is presented in Table 1. Bigger specimens of each species were caught in the Volta estuary than in the Pra estuary.

TABLE 1

Size range of grey mullets caught in the Volta and Pra estuaries

Species	Volta estuary		Pra estuary	
	SL (mm)	Wt (g)	SL (mm)	Wt (g)
<i>L. falcipinnis</i>	38-239	1.1-262.3	42-217	1.3-205.6
<i>L. dumerilii</i>	45-233	1.6-216.3	47-230	2.3-238.7
<i>L. grandisquamis</i>	–	– 70-163	7.9-104.8	
<i>M. bananensis</i>	54-195	3.6-174.5	47-193	2.1-153.2
<i>M. curema</i>	54-270	3.6-444.1	50-158	3.4-78.4
<i>M. cephalus</i>	77-570	10.2-3300	115-355	34.8-1250

- Absent

Composition of stomach contents

Apart from fish scales and eggs that were found in the stomachs of *L. falcipinnis* and an unidentified species of red alga found in a few stomachs of *L. grandisquamis* and *M. bananensis* from the Pra estuary, all the other items were common in the stomachs of all the species from the two estuaries. Twelve species of diatoms, 12 of green algae and 10 species of blue-green algae were identified (Table 2). The most common diatoms found in the stomachs were species of *Navicula*, *Surirella*, *Synedra*, *Gyro-sigma*, *Nitzschia*, and *Cyclo-tella*. The blue-green algae consumed consisted mainly of filamentous types (e.g. *Lyngbya* spp., *Oscilla-toria* spp. and *Anabaena* spp.) and unicellular forms (e.g. *Merismopedia* sp.), while the green algae were mostly species of desmids and filamentous types (e.g. *Chaetophora* sp., and *Ulothrix* sp.). Copepods and protozoans (mainly foraminiferans) were the dominant zooplankton in the diet.

TABLE 2

List of stomach contents of grey mullets from the Volta and Pra estuaries

Stomach contents

Bacteria

Benthic organisms

Polychaetes

Nematodes

Zooplankton

Annelid larvae

Crustacean larvae

Protozoans

Dinoflagellates

Copepods

Ostracods

Rotifers

Cladocerans

Diatoms

Surirella sp.

Gyrosigma sp.

Navicula sp.

Nitzschia sp.

Synedra sp.

Stephanodiscus sp.

Cyclotella sp.

Gomphonema sp.

Pinnularia sp.

Diatoma sp.

Melosira sp.

Cymbella sp.

Sand particles

Detritus

Stomach contents

Blue-green algae

Spirulina sp.

Chroococcus sp.

Lyngbya sp.

Anabaena sp.

Oscillatoria sp.

Merismopedia sp.

Microcystis sp.

Calothrix sp.

Agmellum sp.

Gomphosphaeria sp.

Green algae

Closterium sp.

Pediastrum sp.

Staurastrum sp.

Ankistrodesmus sp.

Schizomeris sp.

Chaetophora sp.

Spirogyra sp.

Ulothrix sp.

Microspora sp.

Prasiola

Stichococcus sp.

Scenedesmus sp.

Unidentified filamentous green algae

Red algae

Fish scales

Fish eggs

Data on stomach contents from the monthly samples were pooled to assess their overall composition in the various species (Fig. 2). This was done after it had been ascertained that there was no appreciable change in the stomach contents throughout the seasons and also among the various size groups of a given species. Bacteria, nematodes, red algae and other organisms as well as fish scales, whose composition in the diet was less than 1.0%, were not included in the analysis. Benthic organisms were, therefore, made up of only polychaetes. Sand particles, diatoms, detritus and green algae were the most important food items, with sand particles forming the bulk of the diet of all the species from both estuaries. Polychaetes also formed an important component of the diet of *M. cephalus* from both estuaries.








Fig. 2. Composition of stomach contents of grey mullets from the Volta and Pra estuaries

Cluster dendograms (Fig. 3), based on frequency of items consumed by the different species in each estuary, show that *L. falcipinnis* and *M. bananensis* frequently consumed similar items in both estuaries. Also, *L. dumerilii* and *L. grandisquamis* ate similar items in the Pra estuary. The items found in the stomachs of *M. curema* in the two estuaries were more related to that found in *L. falcipinnis* and *M. bananensis* in both places and also to that found in the stomachs of *L. dumerilii* and *L. grandisquamis* in the Pra estuary. *M. cephalus* in the two estuaries had little interaction with the other species with respect to the items found in their stomachs. This is, most probably, due to the presence of more polychaetes in their stomachs compared to the other species.



Particle size selection

Species from the Volta estuary generally ingested larger sand particles than their counterparts from the Pra estuary (Table 3). In each estuary, *L. dumerilii* ingested the widest range of sand particles from 41.2 to 1195.8 μm with a mean size of 248.8 μm in the Volta estuary, and from 33.0 to 1649 μm with a mean size of 194.6 μm in the Pra estuary. The most frequently ingested sand particles by *L. dumerilii*, *M. curema* and *M. cephalus* in the Volta estuary

measured 120–139 μm . In the Pra estuary, the most selected particle size by *L. dumerilii* was 80–99 μm , similar to the commonest size ingested by *L. grandisquamis* in the same estuary. The size of sand particles filtered by the latter species, however, ranged from 31.4 to 270.4 μm . The smallest size range of 16.5–255.7 μm was ingested by *M. bananensis* in the Pra estuary while in the Volta estuary, *L. falcipinnis* filtered the smallest size range, 33.0–479.4 μm . The most ingested particle size ranged from 40 to 59 μm for *M. bananensis*, and from 80 to 99 μm for *L. falcipinnis*.

TABLE 3

Size of sand particles ingested by grey mullets from the Volta and Pra estuaries

<i>Species</i>	<i>Estuary</i>	<i>Particle size range (μm)</i>	<i>Modal class (μm)</i>	<i>Mean particle diameter \pm S.E (μm)</i>
<i>L. falcipinnis</i>	Volta	33.0 – 479.4	60-79	126.7 \pm 5.8
	Pra	31.2 – 479.4	40-59	118.3 \pm 6.0
<i>L. dumerilii</i>	Volta	41.2 – 1195.8	120-139	248.8 \pm 13.7
	Pra	33.0 – 1649.4	80-99	194.6 \pm 12.9
<i>L. grandisquamis</i>	Volta			
	Pra	-31.2 – 270.4	-80-99	-99.7 \pm 3.7
<i>M. bananensis</i>	Volta	41.2 – 494.8	80-99	132.9 \pm 6.4
	Pra	16.5 – 255.7	40-59	71.9 \pm 3.5
<i>M. curema</i>	Volta	33.0 – 618.6	120-139	156.5 \pm 6.6
	Pra	31.2 – 527.8	60-79	123.1 \pm 6.1
<i>M. cephalus</i>	Volta	41.2 – 783.5	120-139	213.3 \pm 9.7
	Pra	31.2 – 675.4	100-119	187.3 \pm 7.0

Daily feeding cycle

Changes in the mean index of stomach fullness of the mullets in the Volta estuary over two 24-h periods are illustrated in Fig. 4. The main feeding period in the Volta estuary was during the day between 08.00 h and 12.00 h for all the species. There was little or no feeding during the night, except for *M. cephalus* for which feeding activity peaked at 20.00 h during the first 24-h sampling. The peak in feeding activity for *L. dumerilii* and *L. falcipinnis* occurred at 08.00 h while the peak for *M. bananensis* and *M. curema* occurred at 12.00 h on both sampling occasions. The highest feeding activity on both occasions was shown by *L. dumerilii* which consumed 2.3 and 2.6% of its body weight of food while *M. cephalus* was the least active feeder.




Fig. 4. Tidal pattern and feeding periodicity of grey mullets in the Volta estuary during 11-12 November 1998 (—) and 9-10 June 1999 (---). Vertical bars = \pm SE
Intestine length to standard length ratio (Relative gut length)

The least mean ratio of the length of the intestine to standard length of fish (relative gut length) was 1.82 and 1.72 for *L. dumerilii* in the Volta and Pra estuaries, respectively. *M. cephalus* in the Volta estuary had the highest ratio (4.56) while in the Pra estuary the longest relative gut length of 4.33 was calculated for *L. falcipinnis* (Table 4). The relative gut length of *M. cephalus* in the Pra estuary was not determined because of its low sample size. To compare the relative gut lengths of a given species from the two sites, a Student's *t*-test was performed (Sokal & Rohlf, 1981). The relative gut length of *L. falcipinnis* and *L. dumerilii* in the Volta estuary was significantly different ($p < 0.05$) from specimens in the Pra estuary. However, those of *M. bananensis* and *M. curema* in the Volta estuary were fairly similar to their counterparts in the Pra estuary.

TABLE 4

Relative gut length and standard length of grey mullets from the Volta and Pra estuaries

Species	Number of specimens analysed	Range of standard length (mm)	Range of gut length (mm)	Relative gut length	
				Range	Mean \pm S.E.
<i>Volta estuary</i>					
<i>L. falcipinnis</i>	106	125–232	270–1054	2.00–4.59	3.89 \pm 0.07
<i>L. dumerilii</i>	102	65–200	120–415	1.18–2.57	1.82 \pm 0.03
<i>M. bananensis</i>	78	62–190	162–820	2.61–5.07	3.76 \pm 0.08
<i>M. curema</i>	124	87–235	265–1475	2.48–5.59	3.96 \pm 0.05
<i>M. cephalus</i>	32	113–375	380–1565	3.36–5.27	4.56 \pm 0.10
<i>Pra estuary</i>					
<i>L. falcipinnis</i>	143	70–120	230–580	2.95–5.92	4.33 \pm 0.06
<i>L. dumerilii</i>	92	60–168	88–370	1.36–2.50	1.72 \pm 0.03
<i>L. grandisquamis</i>	63	80–160	105–390	1.08–2.56	2.00 \pm 0.06
<i>M. bananensis</i>	88	75–170	160–600	2.67–4.58	3.81 \pm 0.05
<i>M. curema</i>	104	70–113	170–505	2.46–5.05	4.01 \pm 0.05

Discussion

The diet of all the species in the two estuaries was similar, but slight variations in proportions of food items were noted. The food consisted mostly of diatoms, algae, detrital material and sand particles. Polychaetes were also prominent, especially in the diet of *M. cephalus*. The diet of the mullets examined in this study does not differ appreciably from that of species studied elsewhere in West Africa (Fagade & Olaniyan, 1973; King, 1988; Bruhlet, 1975; Payne, 1976; Albaret & Legendre, 1985) where mullets have been reported to feed mainly on diatoms, organic detritus and sand grains. Although the nutritional value of detritus may be negligible, it is considered a good source of vitamin B₁₂ for mullets (Vallet *et al.*, 1970 quoted in Eggold & Motta, 1992) and is also richly associated with bacteria and protozoa which may be of some nutritional value to the fish (Bruslé, 1981).

Even though sand particles occurred in the stomachs of all the species, they were predominant in the stomachs of *L. dumerilii*, forming 61% and 57% of the stomach contents in the Volta and Pra estuaries, respectively. Ingested sand particles are supposedly helpful in the grinding of food particles in the thick-walled pyloric stomach, which acts as a gizzard (Thomson, 1966). Apart from the 'grinding' action, sand particles with the associated micro-organisms serve as a source of nutrition (Hickling, 1970; Odum, 1970).

It has been suggested that grey mullets ingest sand particles of a selected range (Odum, 1968). This was evident in this study where it was found that the different species in each estuary selected different size ranges of sand particles, with *L. dumerilii* selecting the widest range in both estuaries. The particle size consumed by each species varied according to locality, probably due to differences in the substrate in each estuary. Ingestion of larger particles by *L. dumerilii* compared to other mugilids has been reported by Masson & Marais (1975) from the Zwartkops estuary in Cape Province in South Africa, Payne (1976) from the Black Johnson estuary in Sierra Leone and Blay (1995a) from the Elmina lagoon in Ghana.

Even though *L. dumerilii* ingested the widest range of sand particles, the size it preferred most was also ingested by *M. curema* and *M. cephalus* in the Volta estuary, and by *L. grandisquamis* in the Pra estuary. In the former estuary, *L. dumerilii* showed a preference for zooplankton and diatoms while *M. cephalus* and *M. curema* ingested less zooplankton but more of green algae and polychaetes. In the Pra estuary, *L. grandisquamis* ingested more of diatoms while polychaetes and zooplankton were prominent in the diet of *L. dumerilii*. Thus, species that ingested similar modal sizes of sand particles showed preferences for different food items. This corroborates the findings of Blaber (1976) who noted differences in the food of *L. dumerilii* and *M. cephalus* although they selected similar sand

particles. It would also appear from the results of this study that species that ingested finer particles consumed more plant material and organic matter, while those that ingested coarser particles showed preference for substrates rich in animal material.

According to Nilson (1967), when two closely-related species coexist in one environment, various mechanisms may occur which permit this coexistence. One of these is the partitioning of resources. The preference of each grey mullet species in the two estuaries for a particular particle size was a way of partitioning the resource to avoid interspecific competition, thus ensuring their coexistence.

Apart from detritus and sand particles, results on food habits from the current study confirms the findings of Blay (1995a) who reported on the diet of juveniles of four mullet species (*L. falcipinnis*, *L. dumerilii*, *M. curema* and *M. bananensis*) from the Elmina lagoon in Ghana. Comparatively, mullets in the Elmina lagoon utilize a wider range of food items than was found in the current study. The composition of the diet of grey mullets from different localities is likely to differ, depending on the abundance and types of food organisms present (Blaber, 1977; Bruslé, 1981). The wide range of food items of mugilids has also been reported by Cain & Dean (1976) who explained that such flexibility in feeding ensures a constant energy supply for the sustenance of the population.

The herbivorous feeding habit of adult mullets outside the West African sub-region have been reported by Blaber (1977), De Silva & Wijeyaratne (1977), Whitfield & Blaber (1978), Marais (1980) and Minckley (1982). The ecological importance of such a diet has been recognized by Odum (1970), who considered mullets as important in providing the herbivore link in the food chain in estuaries and shallow coastal waters. Their low position in the food chain is also an asset in pond culture because production of their preferred food items can easily be enhanced in ponds through inexpensive methods of fertilization.

All the species in the Volta estuary showed a marked temporal feeding pattern during a 24-h period with higher feeding activity during the day (08.00–12.00 h) and very little or no feeding activity during the night (20.00–04.00 h). Thus, the species showed a diurnal feeding habit as reported in lagoons at Elmina and Cape Coast in Ghana (Blay, 1995 a,b). Studies by other researchers, e.g. Blabber (1976) on *L. macrolepis* from the St Lucia Lake system in South Africa, De Silva & Wijeyaratne (1977) on *M. cephalus* from the Negombo Lagoon in Sri Lanka and Collins (1981) on *M. cephalus* from Florida showed that these species fed little at night.

Diurnal feeding activity of the mullets in this study was most probably influenced by photoperiodicity, as noted by Keast & Welsh (1968) for many other fishes. Apart from this, other factors, e.g. tides and salinity, have been cited to influence feeding intensity in mugilids (Marais, 1980; Almeida *et al.*, 1993) and Odum (1970) reported a higher feeding intensity in mullets at high tide. No such relationship was, however, found for species in the Elmina lagoon (Blay, 1995a). In the present study, tides apart from photoperiodicity may have influenced the feeding activity of *L. falcipinnis*, *L. dumerilii* and *M. cephalus* with high feeding intensity at low tides. There was, however, no definite relationship between tides and the feeding activity of *M. bananensis* and *M. curema*.

Peak feeding activity of the fishes occurred about the same time in both experimental periods. Differences, however, occurred in the peak feeding time among the species. Different mugilid species have been reported to feed at different times (Marais, 1980; Drake, Arias & Gallegos, 1984). Differences in the peak feeding time among the grey mullet species in the Volta estuary could be one of the probable mechanisms for avoiding competition as suggested by Blaber (1976) for sympatric mullets.

Apart from *L. dumerilii* from both estuaries and *L. grandisquamis* whose relative gut length ranged from 1.72 to 2.0, all the other species had higher relative gut lengths of between 3.7 and 4.5. Results of the present study show that populations with a higher relative gut length tended to consume more plant material, as shown by *L. falcipinnis* and *M. curema* and also reported by Hickling (1970) and Collins (1981). Longer guts could be an adaptation to increasing the digestive area and ensure that such food items are properly digested. The higher consumption of zooplankton by *L. dumerilii* and *L. grandisquamis* may be related to their smaller relative gut length, although diatoms were also major food items.

Conclusion

Stomach contents of grey mullets in the two estuaries consisted of mainly diatoms, detrital material and sand particles. They showed a marked temporal feeding pattern with high feeding activity during the day. The peak of feeding time and the size of sand particles filtered differed among the species. Those found to ingest the same modal size of sand particles showed preference for different food items. These could be strategies adopted to avoid interspecific competition and ensure their coexistence. Their low position in the food chain is an asset in pond culture because production of their preferred food items can easily be enhanced in ponds through inexpensive methods of fertilization.

Acknowledgement

The authors are grateful to the National Agricultural Research Programme (NARP) which provided funding for this study. They are also grateful to Water Research Institute of the Council for Scientific and Industrial Research and the Zoology Department of the University of Cape Coast for providing the necessary logistic support.

References

- Albaret J.-J. and Legendre M.** (1985). Biologie et ecologie des *Mugilidae* en lagune Ébrié (Côte d'Ivoire) Intérêt potentiel pour l'aquaculture lagunaire. *Revue d'Hydrobiologie Tropical* **18**(4): 281–304.
- Almeida P. R., Moreira F., Costa J. L., Assis C. A. & Costa M. J.** (1993). The feeding strategies of *L. ramanda* (Risso, 1826) in fresh and brackish water in the River Tagus, Portugal. *J. Fish Biol.* **42**: 95–107.
- Bardach J. E., Ryther J. H. and McLaren W. O.** (1972). *Aquaculture: The farming and husbandry of freshwater and marine organisms*. New York, Wiley Interscience. 868 pp.
- Blaber S. J. M.** (1987). Factors affecting recruitment and survival of mugilids in estuaries and coastal waters of Southern Africa. *Am. Fish. Soc. Symp.* **1**: 507–518.
- Blaber S. J. M.** (1977). The feeding ecology and relative abundance of mullet (Mugilidae) in Natal and Pondoland estuaries. *Biol. J. Linn. Soc.* **9**: 259–275.
- Blaber S. J. M.** (1976). The food and feeding ecology of Mugilidae in the St. Lucia Lake system. *Biol. J. Linn. Soc.* **8**: 267–227.
- Blay J. Jr.** (1995a). Food and feeding habits of four species of juvenile mullets (Mugilidae) in a tidal lagoon in Ghana. *J. Fish Biol.* **46**: 134–141
- Blay J. Jr.** (1995b). Food habits and diel feeding periodicity of juvenile sicklefin mullet, *Liza falcipinnis* (Pisces: Mugilidae), in a 'closed' tropical lagoon. *Arch. Hydrobiol.* **135** (2): 271–281.
- Bruhlet J.** (1975). Observations on the biology of *Mugil cephalus ashenteensis* and the possibility of its culture on the Mauritanian coast. *Aquaculture* **5**: 271–281.
- Bruslé J.** (1981). Food and feeding in grey mullets. In *Aquaculture of the grey mullets* (O.H. Oren, ed.). Cambridge Univ. Press. pp. 185–217.
- Cain R. L. and Dean J. M.** (1976). Annual occurrence, abundance and diversity of fish in a south Carolina intertidal creek. *Mar. Biol.* **36**: 369–379.
- Collins M. R.** (1981). The feeding periodicity of striped mullet, *Mugil cephalus* L. in two Florida habitats. *J. Fish Biol.* **19**: 307–315.
- De Silva S. S. and Wijeyaratne M. J. S.** (1977). Studies on the biology of young mullet, *Mugil cephalus*, L. II. Food and feeding. *Aquaculture* **12**: 157–167.
- Dickson B. K. and Benneh G.** (1977). *A new geography of Ghana*. London: Longman Group Ltd. 173 pp.
- Drake P. Arias A. M. and Gallegos L.** (1984). Biología de los Mugilidos (Osteichthyes, Mugilidae) en los esteros de las salinas de San Fernanda (Cadiz). III. Hábitos alimentarios y su relación con la morfometría del aparato digestivo. *Investigaciones Pesqueras* **48**: 337–367.
- Eggold B. T. and Motta P. J.** (1992). Ontogenetic dietary shifts and morphological correlates in striped mullet, *Mugil cephalus*. *Envir. Biol. Fishes* **34** (2): 139–158.
- Fagade S. O. and Olaniyan C. I. O.** (1973). The food and feeding interrelationships of the fishes in the Lagos Lagoon. *J. Fish Biol.* **5**: 205–225.
- Ferrari I. and Chiaregato A. R.** (1981). Feeding habits of juvenile stages of *Sparus auratus* L., *Decentrachus labrax* L. and Mugilidae in a brackish embayment of the Po river delta. *Aquaculture* **25** (2–3): 243–257.
- Hickling C. F.** (1970). A contribution to the natural history of the English grey mullets (Pisces, Mugilidae). *J. Mar. Biol. Assoc. U.K.* **50**: 609–633.
- Hureau J. C.** (1966). Biologie comparée de quelques poissons antarctique (Nototheniidae). *Bull. Inst. Oceanogr. Monaco* **68**: 1–244.
- Hynes H. B. N.** (1950). The food of fresh water sticklebacks (*Gasterosteus aculeatus* and *Pygosteus pungitius*) with a review of methods used in studies of the food of fishes. *J. Anim. Ecol.* **19**: 36–58.
- International Centre for Aquatic Resources Management (ICLARM)** (1980). Introduction. In *Review of breeding and propagation techniques for grey mullet, Mugil cephalus* L. (C.E. Nash and Z.H. Shehadeh, ed.). ICLARM Studies and Reviews 3. 87 pp.
- Ikomi R. B.** (1990). On the ecology of the grey mullet (*Mugil cephalus* L.) in Kulama Creek (Niger Delta, Nigeria). *Acta Hydrobiol.* **32** (1/2): 237–249.
- Keast A. and Welsh L.** (1968). Daily feeding periodicities, food uptake rates and dietary changes with hour of day in some lake fishes. *J. Fish. Res. B. Can.* **25**: 1133–1144.
- King R. P.** (1988). New observations on the trophic ecology of *Liza grandisquamis* (Valenciennes, 1836) (Pisces: Mugilidae) in the Bonny river, Niger Delta, Nigeria. *Cybium* **12** (1): 23–36.
- Koutrakis E. T., Sinis A. I. and Economidis P. S.** (1994). Seasonal occurrence, abundance and size distribution of grey mullet fry (Pisces: Mugilidae) in the Porto-Lagos lagoon and Lake Vistonis (Aegean Sea, Greece). *Bamidgeh* **46** (4): 182–196.
- Lévêque C., Paugy D. and Teugels G. G.** (1992). *Faune des poissons d'eaux douces et saumâtres de l'Afrique de l'Ouest (Tome 2)*. Collection Faune Tropical no. XXVIII. ORSTOM et MARC.

- Masson H. and Marais J. F. K.** (1975). Stomach content analyses of mullets from the Zwartkops estuary. *Zoologica Afr.* **10**: 193–207.
- Marais J. F. K.** (1980). Aspects of food intake, food selection and alimentary canal morphology of *Mugil cephalus* L. *Liza tricuspidens* Smith, *L. richardsoni* Smith and *L. dumerili* Steindachner. *J. exp. mar. Biol. Ecol.* **44**: 193–210.
- Minckley W. L.** (1982). Trophic interrelations among introduced fishes in the lower Colorado River, Southeastern U.S., *Calif. Fish Game* **68**: 78–89.
- Nilson N. A.** (1967). Interactive segregation between fish species. In *The Biological basis of freshwater fish production* (S. D. Gerking, ed.), pp. 295–313. Oxford: Blackwell Scientific Publications.
- Obodai E. A., Yankson K. and Blay J. Jr.** (1996). Seasonal changes in hydrographic factors and breeding in two populations of *Crassostrea tulipa* (Lamarck). *Ghana J. Sci.* **31–36** (1991–1996): 45–51.
- Odum W. E.** (1970). Utilisation of the direct grazing plant and detritus food chains by the striped mullet, *Mugil cephalus*. In: *Marine food chains*. (J.H. Steele, ed.), pp. 220–240. London: Oliver and Boyd..
- Odum W. E.** (1968). Ecological significance of fine particle selection by the striped mullet, *Mugil cephalus* L. *Limnol. Oceanogr.* **13**: 92–98.
- Payne A. L.** (1976). The relative abundance and feeding habits of the grey mullet species occurring in an estuary in Sierra Leone, West Africa. *Mar. Biol.* **35**: 277–286.
- Schneider W.** (1990). *Field guide to the commercial marine resources of the Gulf of Guinea*. FAO, Rome. RAFR/FI/90/2.
- Sokal R. R. and Rohlf F. J.** (1981). *Biometry: The principles and practices of statistics in biological research*. San Francisco: W.H. Freeman and Company. 859 pp.
- Thomson J. M.** (1966). The grey mullets. *Oceanogr. Mar. Biol.* **4**: 301–335.
- Whitfield A. K. and Blaber S. J.** (1978). Resource aggregation among illiophagous fish in St. Lucia, Zululand. *Envi. Biol. Fish* **3**: 293–296.
- Wijeyaratne M. J. S. and Costa H. H.** (1986). On the biology of an estuarine population of grey mullet, *Mugil cephalus* L. in Negambo lagoon, Sri Lanka. *Cybium* **10**: 351–363.